



PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM  
Search PubMed for Limits Preview/Index History Clipboard  
About Entrez

Entrez PubMed  
Overview  
Help | FAQ  
Tutorial  
New/Noteworthy

PubMed  
Services  
Journal Browser  
MeSH Browser  
Single Citation  
Matcher  
Batch Citation  
Matcher  
Clinical Queries  
Cubby

Related  
Resources  
Order Documents  
Grateful Med  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

Privacy Policy

1: Exp Neurol 1996 Jul;140(1):1-13

Related Articles, Books, LinkOut



## Xenotransplantation of porcine fetal ventral mesencephalon in a rat model of Parkinson's disease: functional recovery and graft morphology.

Galpern WR, Burns LH, Deacon TW, Dinsmore J, Isacson O.

Neurogeneration Laboratory, McLean Hospital, Harvard Medical School, Massachusetts 02178, USA.

Neurotransplantation of human fetal dopamine (DA) neurons is currently being investigated as a therapeutic modality for Parkinson's disease (PD). However, the practical limitations of human fetal transplantation indicate a need for alternative methodologies. Using the 6-hydroxydopamine rat model of PD, we transplanted dopaminergic neurons derived from Embryonic Day 27 porcine fetuses into the denervated striatum of cyclosporine-A (CyA)-treated or non-CyA-treated rats. Functional recovery was assessed by amphetamine-induced rotation, and graft survival and morphology were analyzed using neuronal and glial immunostaining as well as in situ hybridization with a porcine repeat element DNA probe. A significant, sustained reduction in amphetamine-induced rotational asymmetry was present in the CyA-treated rats whereas the non-CyA-treated rats showed a transient behavioral recovery. The degree of rotational recovery was highly correlated to the number of surviving transplanted porcine dopaminergic neurons. TH+ neuronal survival and graft volume were significantly greater in the CyA-treated group as compared to the non-CyA group. By donor-specific neuronal and glial immunostaining as well as donor-specific DNA labeling, we demonstrate that porcine fetal neuroblasts are able to survive in the adult brain of immunosuppressed rats, mediate functional recovery, and extensively reinnervate the host striatum. These findings suggest that porcine DA neurons may be a suitable alternative to the use of human fetal tissue in neurotransplantation for PD.

PMID: 8682173 [PubMed - indexed for MEDLINE]

Abstract